
Journal of Clinical Pathology



Cover legend: Intravascular platelet aggregates (arrows) in an intratesticular kaposiform hemangioendothelioma. From Costa et al, page 625

Receive regular table of contents by email. Register using this QR code.





This article has been chosen by the Editor to be of special interest or importance and is freely available online



This article has been made freely available OPEN ACCESS online under the BMJ Journals Open Access scheme. See http://jcp.bmj.com/ site/about/guidelines.xhtml#open



This journal is a member of and subscribes to the principles of the Committee on Publication

www.publicationethics.org.uk





Contents

Editorial

Diagnosing vascular invasion in colorectal carcinomas: improving reproducibility and potential pitfalls G Cserni, I Sejben, R Bori

Gene of the month

548 Gene of the month: KRAS R Chetty, D Govender

Original articles

Pathological diagnostic criterion of blood and lymphatic vessel invasion in colorectal cancer: a framework for developing an objective pathological diagnostic system using the Delphi method, from the Pathology Working Group of the Japanese Society for Cancer of the Colon and Rectum M Kojima. H Shimazaki, K Iwaya, M Kage, J Akiba, Y Ohkura, S Horiguchi, K Shomori, R Kushima, Y Ajioka, S Nomura, A Ochiai

Urethral caruncle: a lesion related to IgG4-associated sclerosing disease? S R Williamson, M Scarpelli, A Lopez-Beltran, R Montironi, M R Conces, L Cheng

Strong expression of the neuronal transcription factor FOXP2 is linked to an increased risk of early PSA recurrence in ERG fusion-negative cancers L Stumm. L Burkhardt, S Steurer, R Simon, M Adam, A Becker, G Sauter, S Minner, T Schlomm, H Sirma, U Michl

- Intramuscular corpora amylacea adjacent to ileal low-grade neuroendocrine tumours (typical carcinoids): a light microscopic, immunohistochemical and ultrastructural study [F Hechtman, R E Gordon, N Harpaz
- 573 IGFBP-5 overexpression as a poor prognostic factor in patients with urothelial carcinomas of upper urinary tracts and urinary bladder P-I Liang, Y-H Wang, T-F Wu, W-R Wu, A C Liao, K-H Shen, C-H Hsing, Y-L Shiue, H-Y Huang, H-P Hsu, L-T Chen, C-Y Lin, C Tai, J-Y Wu, C-F Li
- A simplified economic approach to thyroid FNA cytology and surgical intervention in thyroid nodules D N Poller, P Kandaswamv

July 2013 Volume 66 Issue 7

- Frequent expression of follicular dendritic cell markers in Hodgkin lymphoma and anaplastic large cell lymphoma S H Kim, J-Y Choe, Y Jeon, J Huh, HR Iung, Y-D Choi, H-I Kim, HI Cha, WS Park, J E Kim
- Mast cells in infantile haemangioma possess a primitive myeloid phenotype T Itinteang, S T Tan, J Jia, R Steel, E L Laing, H D Brasch, D J Day
- Sarcomatoid carcinoma represents a complete phenotype with various pathways of epithelial mesenchymal transition C O Sung, H Choi, K-W Lee, S-H Kim
- PARP expression in germ cell tumours M Mego. Z Cierna, D Svetlovska, D Macak, K Machalekova, V Miskovska, M Chovanec, V Usakova, J Obertova, P Babal, J Mardiak
- Vasohibin-1 is a new predictor of disease-free survival in operated patients with renal cell carcinoma N Kanomata, Y Sato, Y Miyaji, A Nagai, T Moriya

Short reports

- 620 Vitamin D: a negative acute phase reactant J L Waldron, H L Ashby, M P Cornes, J Bechervaise,
 C Razavi, O L Thomas, S Chugh, S Deshpande, C Ford,
 R Gama
 Intratesticular kaposiform haemangioendothelioma in d
- adults: a report of two cases F D Costa, A L Folpe
- Construction of tissue microarrays using pre-existing ar technologies slides as source of tissue when paraffin blocks are unavailable F-M Deng, Y Zhao, X Kong, P Lee, J Melamed Script 627 Construction of tissue microarrays using pre-existing

PostScript

630 Correspondence

J Clin Pathol: first published as on 1 July 2013. Downloaded from http://jcp.bmj.com/ on April 28, Protected by copyright, including for uses related to text and data mining, Al training, and simila